

ORIGINAL ARTICLE

Laminectomy plus Fusion versus Laminectomy Alone for Lumbar Spondylolisthesis

Zoher Ghogawala, M.D., James Dziura, Ph.D., William E. Butler, M.D.,
Feng Dai, Ph.D., Norma Terrin, Ph.D., Subu N. Magge, M.D.,
Jean-Valery C.E. Coumans, M.D., J. Fred Harrington, M.D.,
Sepideh Amin-Hanjani, M.D., J. Sanford Schwartz, M.D., Volker K.H. Sonntag, M.D.,
Fred G. Barker, II, M.D., and Edward C. Benzel, M.D.

ABSTRACT

BACKGROUND

The comparative effectiveness of performing instrumented (rigid pedicle screws affixed to titanium alloy rods) lumbar spinal fusion in addition to decompressive laminectomy in patients with symptomatic lumbar grade I degenerative spondylolisthesis with spinal stenosis is unknown.

METHODS

In this randomized, controlled trial, we assigned patients, 50 to 80 years of age, who had stable degenerative spondylolisthesis (degree of spondylolisthesis, 3 to 14 mm) and symptomatic lumbar spinal stenosis to undergo either decompressive laminectomy alone (decompression-alone group) or laminectomy with posterolateral instrumented fusion (fusion group). The primary outcome measure was the change in the physical-component summary score of the Medical Outcomes Study 36-Item Short-Form Health Survey (SF-36; range, 0 to 100, with higher scores indicating better quality of life) 2 years after surgery. The secondary outcome measure was the score on the Oswestry Disability Index (range, 0 to 100, with higher scores indicating more disability related to back pain). Patients were followed for 4 years.

RESULTS

A total of 66 patients (mean age, 67 years; 80% women) underwent randomization. The rate of follow-up was 89% at 1 year, 86% at 2 years, and 68% at 4 years. The fusion group had a greater increase in SF-36 physical-component summary scores at 2 years after surgery than did the decompression-alone group (15.2 vs. 9.5, for a difference of 5.7; 95% confidence interval, 0.1 to 11.3; $P=0.046$). The increases in the SF-36 physical-component summary scores in the fusion group remained greater than those in the decompression-alone group at 3 years and at 4 years ($P=0.02$ for both years). With respect to reductions in disability related to back pain, the changes in the Oswestry Disability Index scores at 2 years after surgery did not differ significantly between the study groups (-17.9 in the decompression-alone group and -26.3 in the fusion group, $P=0.06$). More blood loss and longer hospital stays occurred in the fusion group than in the decompression-alone group ($P<0.001$ for both comparisons). The cumulative rate of reoperation was 14% in the fusion group and 34% in the decompression-alone group ($P=0.05$).

CONCLUSIONS

Among patients with degenerative grade I spondylolisthesis, the addition of lumbar spinal fusion to laminectomy was associated with slightly greater but clinically meaningful improvement in overall physical health-related quality of life than laminectomy alone. (Funded by the Jean and David Wallace Foundation and others; SLIP ClinicalTrials.gov number, NCT00109213.)

From the Alan L. and Jacqueline B. Stuart Spine Research Center, the Department of Neurosurgery, Lahey Hospital and Medical Center, Burlington (Z.G., S.N.M.), and the Department of Neurosurgery, Massachusetts General Hospital (W.E.B., J.-V.C.E.C., F.G.B.), and Tufts Clinical and Translational Science Institute, Tufts University School of Medicine (N.T.), Boston — all in Massachusetts; Wallace Trials Center, Greenwich Hospital, Greenwich (Z.G.), and Yale Center for Analytical Sciences, Yale School of Public Health, New Haven (J.D., F.D.) — both in Connecticut; the Department of Neurosurgery, University of New Mexico, Albuquerque (J.F.H.); the Department of Neurosurgery, University of Illinois at Chicago, Chicago (S.A.-H.); Perelman School of Medicine (J.S.S.), Wharton School of Business (J.S.S.), and the Leonard Davis Institute (J.S.S.), University of Pennsylvania, Philadelphia; Barrow Neurosurgical Associates, Barrow Neurological Institute, Phoenix, AZ (V.K.H.S.); and the Center for Spine Health and the Department of Neurosurgery, Cleveland Clinic Foundation, Cleveland (E.C.B.). Address reprint requests to Dr. Ghogawala at the Department of Neurosurgery, Lahey Hospital and Medical Center, 41 Mall Rd., Burlington, MA 01805, or at zoher.ghogawala@lahey.org.

Drs. Barker and Benzel contributed equally to this article.

N Engl J Med 2016;374:1424-34.

DOI: 10.1056/NEJMoa1508788

Copyright © 2016 Massachusetts Medical Society.

THE INCREASED USE OF THE LUMBAR SPINAL fusion procedure in the United States, along with the wide variation in practice, is attracting interest from multiple stakeholders, including patients, physicians, payers, and policymakers. In a report published in 2014, spinal fusion (465,000 hospital-based procedures in 2011) accounted for the highest aggregate hospital costs (\$12.8 billion in 2011) of any surgical procedure performed in U.S. hospitals.¹ The randomized, controlled Spine Patient Outcomes Research Trial (SPORT) showed that surgery was superior to nonoperative care for the management of lumbar degenerative spondylolisthesis.² In SPORT, most patients in the surgical group were treated by means of laminectomy with fusion. Herkowitz et al., in a nonrandomized, prospective, comparative study, found that laminectomy with fusion was superior to laminectomy alone; however, to date, there is no class I evidence that laminectomy plus fusion is superior to laminectomy alone for the treatment of degenerative spondylolisthesis.^{3,4} This lack of evidence complicates efforts to guide and standardize practice, such as through dissemination of clinical practice guidelines. A number of prospective studies with at least 5 years of follow-up after surgery have suggested that lumbar decompression without fusion is associated with excellent outcomes.⁵⁻⁷

The hypothesis tested by the Spinal Laminectomy versus Instrumented Pedicle Screw (SLIP) trial was that lumbar laminectomy with instrumented (rigid pedicle screws affixed to titanium alloy rods) fusion would result in greater improvement than that with laminectomy alone in the primary outcome measure — the change in the physical-component summary score of the Medical Outcomes Study 36-Item Short-Form Health Survey (SF-36; range, 0 to 100, with higher scores indicating better physical health–related quality of life) — at 2 years. Here, we report the results of the primary 2-year outcome of the SLIP trial, as well as the longer-term 3-year and 4-year outcomes.

METHODS

STUDY DESIGN AND OVERSIGHT

In this randomized, controlled trial, patients from five centers were assessed for eligibility during the period from March 2002 through August 2009; the majority (51 patients) were enrolled at one site (for details on the enrollment

statistics, see the Supplementary Appendix, available with the full text of this article at NEJM.org). The primary outcome measure, as specified in the final protocol (available at NEJM.org), was the change in the SF-36 physical-component summary score at 2 years, although the change at both 1 year and 2 years was registered as the primary outcome measure in the initial protocol. The initial trial registration included the score on the Oswestry Disability Index (ODI) as an additional primary outcome measure; however, the final trial protocol, which was prepared in July 2006 (before the completion of the trial or any analyses of trial data), specified the ODI score as a secondary outcome measure. The initial protocol also specified that follow-up would continue through 5 years; however, the analysis was restricted to 4-year follow-up data because of funding limitations and high dropout rates after 4 years of follow-up. The study plan called for the enrollment of 100 patients and the random assignment of at least 64 patients, with funding to enroll approximately 40 patients in a parallel registry — an observation cohort for patients who declined to undergo randomization.

Data were managed at the Wallace Clinical Trials Center in Greenwich, Connecticut. There was no industry funding or any other industry involvement in the SLIP trial. The authors vouch for the accuracy and completeness of the data and analyses and for the fidelity of this report to the trial protocol. Institutional review board approval was obtained at all five participating sites. Written informed consent was obtained from all enrolled patients.

PATIENTS

All patients with grade I lumbar spondylolisthesis (degree of spondylolisthesis, 3 to 14 mm) with lumbar stenosis and neurogenic claudication with or without lumbar radiculopathy were eligible for inclusion. Patients were excluded if radiography revealed lumbar instability (motion of >3 mm at the level of listhesis, as measured on flexion–extension radiographs of the lumbar spine), if they were judged by the enrolling surgeon to have lumbar instability because of a history of mechanical low back pain with axial loading of the spine, if they had had previous lumbar spinal surgery, or if they had American Society of Anesthesiologists (ASA) class IV or higher disease (with classes ranging from I to VI and higher classes indicating more severe systemic disease).

Patients were screened and enrolled by trial coordinators at each site. A panel of 10 expert spine surgeons was formed to review a brief clinical vignette plus four standardized radiographic and magnetic resonance images for each patient to assess suitability for randomization. This novel approach appeared to increase patient consent to undergo randomization (see the Supplementary Appendix).^{8,9} Radiographic and magnetic resonance images from each patient were reviewed centrally by two neuroradiologists and one neurosurgeon to verify degenerative lumbar canal stenosis with spondylolisthesis without disk herniation. In addition, independent radiologic review of postoperative computed tomographic scans confirmed adherence to the study protocol.

INTERVENTIONS

All patients underwent either decompression alone (decompression-alone group) or decompression with posterolateral instrumented fusion (fusion group) at the single level of spondylolisthesis. Decompression was performed by means of a complete laminectomy with partial removal of the medial facet joint.^{10,11} Patients in the fusion group underwent a lumbar laminectomy as well as implantation of pedicle screws and titanium alloy rods across the level of listhesis, with a bone graft harvested from the iliac crest.^{12,13} The SLIP trial did not include the use of bone morphogenetic protein, interbody devices, or minimally invasive techniques for the placement of percutaneous pedicle screws. All the surgeons routinely performed both operations tested in the trial; each of the surgeons had performed at least 100 laminectomies and 100 posterolateral fusions for lumbar spondylolisthesis before joining the SLIP trial.

OUTCOME MEASURES

The primary outcome measure was the change in the SF-36 physical-component summary score at 2 years after surgery. The minimal clinically important difference, which was determined on the basis of previous studies, was prespecified to be 5 points.^{14,15} The secondary outcome measure was the change in the disease-specific ODI score (range, 0 to 100, with higher scores indicating more disability related to back pain).¹⁶ The minimal clinically important difference for the ODI was 10 points.^{15,17,18} Initial clinical assessments were performed during routine outpatient visits

at 1.5 months and 3 months by an independent study coordinator who was not aware of the study hypothesis. After 3 months, validated outcome assessment tools (SF-36 and ODI) were mailed to each patient, who then completed and returned them. A study coordinator attempted to contact patients at least three times to improve patient retention. Additional outcome measures that were prespecified in the protocol included operative complications and reoperations. Reoperation was performed at the discretion of the surgeon; patients were contacted annually by independent study coordinators for assessment of the outcomes of reoperation. A prespecified hospital cost analysis was also described in the protocol, although it has not yet been conducted. Although not explicitly described in the protocol, we collected and reported data on estimated blood loss, operative time, and length of stay in the hospital.

STATISTICAL ANALYSIS

The sample size was estimated on the basis of a previous prospective pilot study that was performed by the principal investigator in 2004.¹⁰ No data from the pilot study were included in this report. We assumed a standard deviation of 10 for the change in SF-36 physical-component summary score and a 10% rate of loss to follow-up at 2 years. We estimated that with a sample size of 64 patients (32 patients in each randomized group), the study would have 80% power to detect a between-group difference of 7.5 points in the degree of improvement in SF-36 physical-component summary scores, at a two-sided significance level of 0.05.

The strategy for analysis was developed after the trial was completed but before the examination of the data (see the statistical analysis plan, which is available with the protocol). The baseline characteristics of the patients were compared between the groups with the use of independent-sample t-tests for continuous variables, which are presented as means and standard deviations, and the chi-square test or Fisher's exact test for categorical variables, which are presented as numbers and percentages. Analyses of the primary outcome were performed among all patients who had follow-up assessments, according to their original randomized treatment assignments.

The between-group comparisons of changes

in SF-36 and ODI scores from baseline were made with the use of mixed-effects models for repeated measures. An unstructured covariance matrix was specified to account for the within-patient correlations of repeatedly measured outcomes. Fixed effects for site, treatment (decompression with fusion vs. decompression alone), time (1.5 months, 3 months, 6 months, 1 year, 2 years, 3 years, and 4 years after randomization), and time-by-treatment interaction were included and were reported as least-squares means and 95% confidence intervals. We computed the robust standard errors and test statistics involving the fixed effects by specifying the EMPIRICAL option within the PROC MIXED procedure (SAS Institute). Comparisons of least-squares means between the treatment groups at each time point and between time points within each treatment group were performed with the use of appropriate contrasts within the mixed-effect models for repeated measures.¹⁹

We compared the percentage of patients in the two groups who had a prespecified minimal clinically important difference of 5 points in the SF-36 physical-component summary score^{14,15} at 2 years by fitting a random-intercept logistic-regression model using PROC GLIMMIX (SAS Institute), with adjustment made for the same list of fixed effects. All analyses were performed with SAS software, version 9.4 (SAS Institute).²⁰

RESULTS

PATIENTS

Figure 1 shows the enrollment, randomization, and follow-up for the SLIP trial. Overall, 130 patients were screened and identified as eligible; 24 eligible patients declined to participate in either treatment group. Among the remaining 106 patients, 66 consented to undergo randomization and 40 declined to undergo randomization because they had a clear preference for a particular surgical strategy, although they agreed to remain in an observation group. One patient who was randomly assigned to the decompression-alone group never had surgery. There were no crossovers from either randomized strategy. A total of 14 patients in the two groups underwent a subsequent reoperation; data from these patients were not censored and were included in the analysis of the primary outcome measure

according to the treatment group to which the patient had been randomly assigned.

The mean age of the study population was 67 years, and 80% were women; the age and the preponderance of women are both consistent with findings in previous reports on patients with degenerative spondylolisthesis.² Baseline characteristics of patients randomly assigned to the decompression-alone group and of those assigned to the fusion group are shown in Table 1. The patients in the fusion group had a mean baseline SF-36 physical-component summary score that was 3.2 points lower than that in the decompression-alone group ($P=0.08$). Three of the 28 patients in the fusion group for whom data on ASA classification were available (11%), as compared with none of the 33 in the decompression-alone group with available data, had ASA class III disease ($P=0.09$). The degree of spondylolisthesis was 5.6 mm in the fusion group and 6.5 mm in the decompression-alone group ($P=0.10$).

PRIMARY OUTCOME MEASURE

At 2 years after surgery, patients in the fusion group had a significantly greater increase in the SF-36 physical-component summary score than did those in the decompression-alone group (15.2 points; 95% confidence interval [CI], 10.9 to 19.5; vs. 9.5 points; 95% CI, 5.2 to 13.8) (Table 2). There was a significant between-group difference in the mean treatment effect (i.e., change in SF-36 physical-component summary score from baseline) of 5.7 points (95% CI, 0.1 to 11.3; $P=0.046$). The magnitude of the difference in treatment effect was sustained longitudinally over the 4 years after surgery (difference at 4 years, 6.7 points; 95% CI, 1.2 to 12.3; $P=0.02$) (Table 2 and Fig. 2).

Among the patients who were available for the 2-year follow-up, 24 of 28 in the fusion group and 20 of 29 in the decompression-alone group had a prespecified minimal clinically important difference of 5 points in the SF-36 physical-component summary score. According to a random-intercept logistic-regression model, the predicted rate of a minimal clinically important difference of 5 points at the 2-year follow-up was 91.9% (95% CI, 73.1 to 97.9) among patients in the fusion group and 76.1% (95% CI, 49.7 to 91.1) among patients in the decompression-alone group (difference, 15.8 percentage points; 95% CI, -16.0 to 47.6; $P=0.18$).

No prespecified plan was outlined for the




Table 1. Baseline Characteristics of the Patients.*

Characteristic	Decompression- Alone Group (N=35)	Fusion Group (N=31)
Age — yr	66.5±8.0	66.7±7.2
Female sex — no. (%)	27 (77)	26 (84)
Walking capacity — min†	16.0±14.7	15.1±8.7
Motor deficit — no./total no. (%)‡	8/34 (24)	8/31 (26)
SF-36 physical-component summary score§	34.7±7.0	31.5±7.3
Physical function	44.7±22.8	38.7±23.7
Bodily pain	33.9±16.3	28.1±13.3
Vitality	45.0±19.8	47.7±23.3
ODI score¶	36.3±15.2	38.8±16.0
Body-mass index	27.1±6.1	29.5±6.5
Degree of spondylolisthesis — mm**	6.5±2.3	5.6±2.2
Spinal canal area at the level of listhesis — mm ² **	39.2±45.1	38.1±27.7
Flexion–extension movement at listhesis — mm**	1.3±1.0	1.6±1.4
ASA class III — no./total no. (%)††	0/33	3/28 (11)
Rotatory scoliosis >15 degrees — no./total no. (%)**	3/31 (10)	4/27 (15)
Lateral scoliosis >15 degrees — no./total no. (%)**	7/31 (23)	3/27 (11)
Disk-space height — mm**	7.3±2.6	8.2±2.9

* Plus–minus values are means ±SD. No between-group differences were significant at P<0.05. Three baseline variables differed between the groups with P values between 0.05 and 0.10: the physical-component summary score of the Medical Outcomes Study 36-Item Short-Form Health Survey (SF-36) (P=0.08), American Society of Anesthesiologists (ASA) class (P=0.09), and degree of spondylolisthesis (P=0.10).

† Walking capacity measured the number of minutes a patient could walk without stopping. Data were available for 26 patients in the decompression-alone group and 28 in the fusion group.

‡ Motor deficit refers to measurable weakness in motor function detected on physical examination of a patient.

§ The physical-component summary of the SF-36 includes general health and vitality, physical functioning, role–physical, and bodily pain; scores range from 0 to 100, with higher scores indicating better quality of life.

¶ Scores from the Oswestry Disability Index (ODI) range from 0 to 100, with higher scores indicating more disability related to back pain.

|| The body-mass index is the weight in kilograms divided by the square of the height in meters. Data were available for 33 patients in the decompression-alone group and 28 in the fusion group.

** Values were determined by central independent image review. The degree of spondylolisthesis and disk-space height were measured from a lateral-view lumbar radiograph, obtained with the patient in a standing position. Data on the degree of spondylolisthesis and on disk-space height were available for 35 patients in the decompression-alone group and 29 in the fusion group. Spinal canal area at the level of listhesis was calculated from axial T₂-weighted magnetic resonance images by measurement of the anterior–posterior and lateral dimensions of the thecal sac, as described previously²¹; data were available for 35 patients in the decompression-alone group and 28 in the fusion group. Flexion–extension movement was measured from flexion–extension radiographs at the level of listhesis, obtained with the patient in a standing position; data were available for 32 patients in the decompression-alone group and 28 in the fusion group. Rotatory and lateral scoliosis greater than 15 degrees were assessed from reconstructed lumbar computed tomographic scans.

†† The physical status classification system of the ASA ranges from I to VI, with higher classes indicating greater risk (class I, healthy patients; class II, mild systemic disease; and class III, severe systemic disease limiting normal activity; no patients with ASA class IV [severe systemic disease that is a constant threat to life], class V [moribund patient not expected to survive 24 hours with or without operation], or class VI [brain-dead patient whose organs are being removed for donor purposes] were included in the study).

adjustment of baseline differences between the groups. For three of the baseline values (SF-36 physical-component summary, ASA class, and degree of spondylolisthesis), there were marginal, nonsignificant differences between the treatment groups, with P values between 0.05

Table 2. Changes in SF-36 Physical-Component Summary and ODI Scores from Baseline.*

Outcome	Change from Baseline		Difference in Change, Fusion vs. Decompression-Alone (95% CI)	P Value
	Decompression-Alone Group	Fusion Group		
SF-36 physical-component summary score				
Baseline†	34.7	31.5	NA	NA
1.5 mo	6.3	5.1	-1.1 (-5.9 to 3.7)	0.64
3 mo	7.7	12.2	4.5 (-0.7 to 9.7)	0.09
6 mo	9.2	15.6	6.4 (1.1 to 11.7)	0.02
1 yr	11.3	15.3	3.9 (-1.5 to 9.4)	0.16
2 yr	9.5	15.2	5.7 (0.1 to 11.3)	0.046
3 yr	7.9	15.3	7.4 (1.1 to 13.7)	0.02
4 yr	7.4	14.1	6.7 (1.2 to 12.3)	0.02
ODI score				
Baseline†	36.3	38.8	NA	NA
1.5 mo	-15.3	-12.4	2.9 (-7.4 to 13.2)	0.58
3 mo	-17.0	-22.2	-5.2 (-13.9 to 3.5)	0.24
6 mo	-20.3	-25.9	-5.6 (-14.4 to 3.2)	0.21
1 yr	-22.2	-26.1	-3.9 (-12.9 to 5.0)	0.38
2 yr	-17.9	-26.3	-8.5 (-17.5 to 0.5)	0.06
3 yr	-17.2	-21.8	-4.6 (-14.7 to 5.6)	0.37
4 yr	-14.7	-23.7	-9.0 (-18.0 to 0.1)	0.05

* Data are presented as least-squares mean values of changes in SF-36 physical-component summary scores and ODI scores from baseline at each follow-up point. Adjustment for multiplicity was not applied. NA denotes not applicable.

† The baseline scores shown are the mean values in the group.

and 0.10 (Table 1). When a separate analysis was performed to adjust for these baseline differences, the fusion group had greater (although not significantly greater) increases in SF-36 physical-component summary scores than did the decompression-alone group at 2 years (3.9 points; 95% CI, -1.9 to 9.7; $P=0.19$), 3 years (5.8 points; 95% CI, -0.4 to 12.1; $P=0.07$), and 4 years (4.6 points; 95% CI, -0.9 to 10.2; $P=0.10$).

SECONDARY OUTCOME MEASURES AND SURGICAL COMPLICATIONS

The differences between the treatment groups in the amelioration of disability related to low back pain, as measured by the change in ODI score at 2, 3, and 4 years, were not significant (Table 2). As shown in Figure 2C, the fusion group had a lower rate of reoperation over the course of 4 years than did the decompression-alone group (14% vs. 34%, $P=0.05$). All the reoperations per-

formed in the decompression-alone group were at the index level to address subsequent clinical instability. In contrast, all the reoperations performed in the fusion group were at an adjacent lumbar level (either disk herniation or clinical instability). Obesity (body-mass index [the weight in kilograms divided by the square of the height in meters] >30) was not a risk factor for reoperation (rates of reoperation were 21% among obese patients and 28% among nonobese patients, $P=0.58$). With respect to surgical complications, blood loss, length of stay, and length of procedure were significantly greater in the fusion group than in the decompression-alone group (Table 3).

DISCUSSION


The comparative effectiveness of instrumented fusion with laminectomy versus laminectomy

Figure 2. Quality-of-Life and Disability Scores and Risk of Reoperation.

Shown are unadjusted physical-component summary scores on the Medical Outcomes Study 36-Item Short-Form Health Survey (SF-36; scores range from 0 to 100, with higher scores indicating better quality of life) (Panel A) and scores on the Oswestry Disability Index (scores range from 0 to 100, with higher scores indicating more disability related to back pain) (Panel B), before and after surgery, among patients who were randomly assigned to undergo laminectomy with fusion (fusion group) or decompressive laminectomy alone (decompression-alone group). I bars represent standard errors. The cumulative risk of reoperation over time (Panel C) is shown for all patients in the decompression-alone group and the fusion group who were available for analysis at 3, 6, 12, 24, 36, and 48 months. Data for patients who were lost to follow-up or who died were censored and are represented with a plus sign. The cumulative risk of reoperation was 34% in the decompression-alone group and 14% in the fusion group ($P=0.05$).

alone was the clinical issue examined in the SLIP trial, which screened 130 patients for eligibility, enrolled 106 patients, and randomly assigned 66 patients. The primary outcome measure was the change in physical health–related quality of life at 2 years, as measured by the SF-36 physical-component summary score. Although the outcomes did not differ significantly between the treatment groups at 1 year after surgery, the addition of lumbar fusion to laminectomy was associated with significantly greater increases in the SF-36 physical-component summary score at 2, 3, and 4 years after surgery, which suggests a sustained difference between treatments over time. The between-group differences in the increases in SF-36 physical-component summary score were small but clinically meaningful. We did not observe significant between-group differences with respect to reductions in the ODI score, which was the secondary outcome measure of disability related to back pain.

It is generally agreed that mobile degenerative spondylolisthesis with mechanical low back pain causes instability in the lumbar spine and should be treated with decompression plus fusion.²² One question addressed by the SLIP trial was whether a lumbar laminectomy destabilizes the lumbar spine in the context of a nonmobile degenerative spondylolisthesis. In the SLIP trial, degenerative spondylolisthesis after lumbar laminectomy was



sufficiently unstable to require reoperation in at least one third of the patients. This rate of reoperation after laminectomy was higher than that reported in other studies but was consistent with the rate of 28% reported in administrative data

Table 3. Surgical Complications.*

Variable	Decompression- Alone Group (N=35)	Fusion Group (N=31)	P Value
Estimated blood loss			
No. of patients with data	34	31	
Mean — ml	83.4±63.5	513.7±334.4	<0.001
Length of stay in the hospital			
No. of patients with data	33	30	
Mean — days	2.6±0.9	4.2±0.9	<0.001
Duration of operation			
No. of patients with data	34	30	
Mean — min	124.4±34.2	289.6±66.3	<0.001
Major complications — no./total (%)	2/35 (6)†	1/31 (3)†	1.0

* Plus-minus values are means ±SD.

† The complications in the decompression-alone group included wound infection and new neurologic deficit. The complication in the fusion group was pneumonia. All complications were identified within 30 days. Minor complications were not recorded.

from the state of Washington.²³ Larger prospective registry studies might assess the generalizability of the rate of reoperation in this trial. A newer — but relatively untested — less-invasive strategy of unilateral laminotomy with bilateral decompression may offer an advantage over traditional laminectomy because the midline ligamentous structures are preserved, possibly reducing the risk of reoperations for instability.²⁴ We have previously reported risk factors for instability after laminectomy for degenerative spondylolisthesis that include disk height, facet angle, and motion on flexion–extension radiographs.¹¹ Identifying patients whose spines would be likely to remain stable after surgical decompression may reduce the use of the lumbar spinal fusion procedure and may reduce the rate of complications. We observed a reoperation rate of 14% after laminectomy plus fusion, a rate that was similar to that in SPORT (11%)² and in a recent study by Brodke et al. (13.3%).²⁵

Although SF-36 physical-component summary scores in the fusion group were statistically higher than the scores in the decompression-alone group at 2, 3, and 4 years, the magnitude of the between-group difference was small. Although patients in the decompression-alone group had a significantly higher risk of early reoperation for instability, revision fusion surgery was

associated with subsequent better outcome scores than those before the reoperation. Lumbar fusion was significantly associated with more blood loss and longer operative times and therefore might not be appropriate for elderly patients or for patients with certain coexisting conditions, including osteoporosis.

In the SLIP trial, the strategy for performing a fusion included implantation of rigid pedicle screws affixed to titanium alloy rods, with bone graft harvested from the iliac crest. Currently, some surgeons use minimally invasive techniques and use bone-graft extenders or bone morphogenetic protein instead of bone grafts harvested from the iliac crest. In addition, the use of interbody fusion techniques has increased since the SLIP trial was completed.²⁶ The SPORT spondylolisthesis trial did not identify any one fusion technique as superior to the others²⁷; autografts from the iliac crest were used in nearly one third of the cases, and there were no significant differences in the rate of bony fusion or in the rate of reoperation.²⁸ The most effective method for creating lumbar fusion is not known.

There were marginal, nonsignificant differences in the baseline variables between the patients in the decompression-alone group and those in the fusion group. Some of the differences in the observed outcomes might be attributed to baseline differences rather than to the randomized treatment. Additional studies are important to validate the observations made in the SLIP trial, which might not be generalizable.

The SF-36 physical-component summary has been shown to be a valid, responsive, and reliable tool for the assessment of degenerative lumbar spinal conditions.¹⁷ The assumption we made when calculating the initial sample size estimate, that 10% of the randomly assigned patients would be lost to follow-up, was reasonably accurate at 1 year (11%) and at 2 years (14%); however, by 4 years, 30% of the patients in the initial randomized treatment groups were lost to follow-up. The interpretation of the differences observed at the 3-year and 4-year time points are weakened by the lower rates of follow-up. Future studies will benefit from larger sample sizes that also include valid disease-specific assessments as primary outcomes.

The well-established higher hospital costs of lumbar fusion may suggest that an overall value assessment might favor decompression alone, as

highlighted in a previous study, as well as in another study in this issue of the *Journal* on the comparative effectiveness of adding fusion to decompression for lumbar spinal stenosis.^{29,30} Future economic analyses would probably include loss of productivity, reoperations, and the use of outpatient health resources to compare these surgical approaches over a longer period.³¹

In conclusion, we found that lumbar laminectomy plus fusion was associated with a slightly greater but clinically meaningful improvement in physical health–related quality of life than was laminectomy alone at 2, 3, and 4 years after surgery.

REFERENCES

- Weiss AJ, Elixhauser A, Andrews RM. Characteristics of operating room procedures in U.S. hospitals, 2011. HCUP statistical brief #170. February 2014 (<http://www.hcup-us.ahrq.gov/reports/statbriefs/sb170-Operating-Room-Procedures-United-States-2011.pdf>).
- Weinstein JN, Lurie JD, Tosteson TD, et al. Surgical versus nonsurgical treatment for lumbar degenerative spondylolisthesis. *N Engl J Med* 2007;356:2257-70.
- Resnick DK, Watters WC III, Sharan A, et al. Guideline update for the performance of fusion procedures for degenerative disease of the lumbar spine — part 9: lumbar fusion for stenosis with spondylolisthesis. *J Neurosurg Spine* 2014;21:54-61.
- Herkowitz HN, Kurz LT. Degenerative lumbar spondylolisthesis with spinal stenosis: a prospective study comparing decompression with decompression and intertransverse process arthrodesis. *J Bone Joint Surg Am* 1991;73:802-8.
- Chang HS, Fujisawa N, Tsuchiya T, Oya S, Matsui T. Degenerative spondylolisthesis does not affect the outcome of unilateral laminotomy with bilateral decompression in patients with lumbar stenosis. *Spine (Phila Pa 1976)* 2014;39:400-8.
- Rampersaud YR, Fisher C, Yee A, et al. Health-related quality of life following decompression compared to decompression and fusion for degenerative lumbar spondylolisthesis: a Canadian multicentre study. *Can J Surg* 2014;57(4):E126-33.
- Kelleher MO, Timlin M, Persaud O, Rampersaud YR. Success and failure of minimally invasive decompression for focal lumbar spinal stenosis in patients with and without deformity. *Spine (Phila Pa 1976)* 2010;35(19):E981-7.
- Ghogawala Z, Benzel EC, Heary RF, et al. Cervical spondylotic myelopathy surgical trial: randomized, controlled trial design and rationale. *Neurosurgery* 2014;75:334-46.
- Ghogawala Z, Schwartz JS, Benzel EC, et al. Increased patient enrollment to a randomized surgical trial through equipoise polling of an expert surgeon panel. *Ann Surg* 2015 October 22 (Epub ahead of print).
- Ghogawala Z, Benzel EC, Amin-Hanjani S, et al. Prospective outcomes evaluation after decompression with or without instrumented fusion for lumbar stenosis and degenerative Grade I spondylolisthesis. *J Neurosurg Spine* 2004;1:267-72.
- Blumenthal C, Curran J, Benzel EC, et al. Radiographic predictors of delayed instability following decompression without fusion for degenerative grade I lumbar spondylolisthesis. *J Neurosurg Spine* 2013;18:340-6.
- Kurz LT, Garfin SR, Booth RE Jr. Harvesting autogenous iliac bone grafts: a review of complications and techniques. *Spine (Phila Pa 1976)* 1989;14:1324-31.
- Dimar JR, Glassman SD, Burkus KJ, Carreon LY. Clinical outcomes and fusion success at 2 years of single-level instrumented posterolateral fusions with recombinant human bone morphogenetic protein-2/compression resistant matrix versus iliac crest bone graft. *Spine (Phila Pa 1976)* 2006;31:2534-9.
- Ware J, Kosinski M, Keller D. SF-36 Physical and Mental Health Summary Scales: a manual for users of version 1. 2nd ed. Lincoln, RI: QualityMetric, 2001.
- Copay AG, Glassman SD, Subach BR, Berven S, Schuler TC, Carreon LY. Minimum clinically important difference in lumbar spine surgery patients: a choice of methods using the Oswestry Disability Index, Medical Outcomes Study questionnaire Short Form 36, and pain scales. *Spine J* 2008;8:968-74.
- Fairbank JC, Couper J, Davies JB, O'Brien JP. The Oswestry low back pain disability questionnaire. *Physiotherapy* 1980;66:271-3.
- Ghogawala Z, Resnick DK, Watters WC III, et al. Guideline update for the performance of fusion procedures for degenerative disease of the lumbar spine — part 2: assessment of functional outcome following lumbar fusion. *J Neurosurg Spine* 2014;21:7-13.
- Hagg O, Fritzell P, Nordwall A. The clinical importance of changes in outcome scores after treatment for chronic low back pain. *Eur Spine J* 2003;12:12-20.
- Hedeker D, Gibbons R. Longitudinal data analysis. New York: Wiley, 2006.
- SAS, version 9.4. Cary, NC: SAS Institute, 2013.
- Hamanishi C, Matukura N, Fujita M, Tomihara M, Tanaka S. Cross-sectional area of the stenotic lumbar dural tube measured from the transverse views of magnetic resonance imaging. *J Spinal Disord* 1994;7:388-93.
- Schroeder GD, Kepler CK, Kurd MF, et al. Rationale for the surgical treatment of lumbar degenerative spondylolisthesis. *Spine (Phila Pa 1976)* 2015;40(21):E1161-6.
- Martin BI, Mirza SK, Comstock BA, Gray DT, Kreuter W, Deyo RA. Reoperation rates following lumbar spine surgery and the influence of spinal fusion procedures. *Spine (Phila Pa 1976)* 2007;32:382-7.
- Alimi M, Hofstetter CP, Pyo SY, Paulo D, Härtl R. Minimally invasive laminectomy for lumbar spinal stenosis in patients with and without preoperative spondylolisthesis: clinical outcome and reoperation rates. *J Neurosurg Spine* 2015;22:339-52.
- Brodke DS, Annis P, Lawrence BD, Woodbury AM, Daubs MD. Reoperation and revision rates of 3 surgical treatment methods for lumbar stenosis associated with degenerative scoliosis and spondylolisthesis. *Spine (Phila Pa 1976)* 2013;38:2287-94.
- Kepler CK, Vaccaro AR, Hilibrand AS, et al. National trends in the use of fusion techniques to treat degenerative spondylolisthesis. *Spine (Phila Pa 1976)* 2014;39:1584-9.
- Abdu WA, Lurie JD, Spratt KF, et al.

- Degenerative spondylolisthesis: does fusion method influence outcome? Four-year results of the Spine Patient Outcomes Research Trial. *Spine (Phila Pa 1976)* 2009; 34:2351-60.
28. Radcliff K, Hwang R, Hilibrand A, et al. The effect of iliac crest autograft on the outcome of fusion in the setting of degenerative spondylolisthesis: a subgroup analysis of the Spine Patient Outcomes Research Trial (SPORT). *J Bone Joint Surg Am* 2012;94:1685-92.
29. Kim S, Mortaz Hedjri S, Coyte PC, Rampersaud YR. Cost-utility of lumbar decompression with or without fusion for patients with symptomatic degenerative lumbar spondylolisthesis. *Spine J* 2012;12:44-54.
30. Försth P, Ólafsson G, Carlsson T, et al. A randomized, controlled trial of fusion surgery for lumbar spinal stenosis. *N Engl J Med* 2016;374:1413-23.
31. Soegaard R, Bünger CE, Christiansen T, Høy K, Eiskjaer SP, Christensen FB. Circumferential fusion is dominant over posterolateral fusion in a long-term perspective: cost-utility evaluation of a randomized controlled trial in severe, chronic low back pain. *Spine (Phila Pa 1976)* 2007;32:2405-14.

Copyright © 2016 Massachusetts Medical Society.